



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

Southern Pacific Railroad (here running midway between the Sierra and the lake), the earliest border of that basin, distant quite fourteen miles from the lake-shore as recorded on the maps, but which itself is now several miles inland from the water's edge. The railroad and Tulare City lie on a belt of sandy land, obviously somewhat higher in level than the 'black lands,' and here about eight miles wide. In crossing this belt to the eastward we traverse several bands of 'alkali land,' characterized by the dense growth of *Brizopyrum*, or 'alkali-grass,' and evidently forming a summit plateau on the divide between the Tulare basin proper and the extreme southern branch of the delta bayous¹ of the Kaweah River, called 'Outside Creek,' or Elk Bayou. Approaching this water-course, we again come to a 'black-lands' belt, about three miles wide, which borders the bayou on both sides, and evidently represents an estuary of the time when Tulare Lake was much higher than now, and the Kaweah delta bayous were mere swamps. Another 'alkali-land' belt is crossed after traversing the 'black-lands' of Outside Creek, towards the foot-hills: beyond these, lies a narrow sandy belt corresponding to that along the railroad, as above described. Then, at a distance of some eight miles from the foot-hills, the color of the soil begins to change toward the well-known red tint of the soils resulting from the decomposition of the 'bedrock' slates of the foot-hills; but the ascent is so gradual that to the eye the plain appears as level as ever, although the presence of the ferric hydrate in the soil proves that these 'red lands' were never submerged for any great length of time, since otherwise their iron would have been reduced and leached out, or gathered into 'black gravel' (bog-ore), as is the case in the 'black lands' of the lake and bayou borders. On inquiry, it was learned that all the larger streams of the region (including Tule River and Deer Creek, outside of the Kaweah delta) are accompanied by belts of such 'black land.'

Besides these main bayous, there appear in the sandy lands a few obvious sandy channels, usually dry, but carrying water in time of flood.

But a curious and at present very striking demonstration of the ancient drainage system of the region may be seen in the grain-fields. In consequence of the failure of the usual April rains, most of the wheat-fields of the region are now in a very precarious condition where not irrigated, and much of the wheat sown will not even make hay. Its condition is best on the 'black lands,' and in certain portions of the sandy belt that do not show any obvious difference in soil from adjoining tracts in which the crop is already dried up without having been able to form grain in the ears. The eye quickly recognizes the extraordinary resemblance of the outlines of the dead portions to meandering water-channels, but no difference of surface-level remains to indicate the fact. But, by digging in any part of these meandering belts of desolation, we find the sandy soil becoming sandier as we descend, until finally, at about three feet depth, an almost pure, coarse sand underlies, which obviously cannot raise moisture within reach of the root system. On the adjacent land, where the wheat is still green and growing, we find at the same depth a subsoil of increased closeness and capillary power, which keeps the moisture below within reach of the roots.

Thus a bird's-eye-view photograph taken of this region now, would show, traced out in minute detail by the color-contrast between the living and the dead grain, the ancient drainage of the country, of which its surface at present shows no indication, together with the broad bands of the ancient estuaries that have formed the 'black lands,' characterized by green, growing grain or an extraordinarily luxuriant growth of oaks, that likewise outlines the ancient margin of Lake Tulare.

We thus obtain a chapter of the geological history of the valley from a mere reconnaissance such as any one desiring to invest in its lands would need to make. The significance of the 'alkali lands' in both points of view remains for a future discussion.

It is hardly necessary to dwell upon the interest attaching to the study of these features, whether from a practical or a purely scientific standpoint.

¹ It should be understood that the Kaweah River, emerging from a cañon of the Sierra next to southward of King's River, divides into a number of forks or bayous immediately upon entering the valley plain. The extensive delta region thus formed is one of the richest, as well as the only forest-grown area of the great valley of California.

One point, however, should be specially noted; namely, that a great many of the characteristic marks of these late quaternary events are rapidly disappearing before the advance of cultivation, and the replacement of the native plant-growth (the result of secular co-adaptation of soils and plants) and of the natural surface by the well-known results of agricultural operations. The latter are already obliterating, on large tracts, the singular 'hog-wallow' mounds that form so striking a feature and so difficult a problem, the solution of which must largely depend upon the geographical distribution of these swarms of mammillary elevations.

It is hard to see on what ground the study of these latest phenomena, connecting the present with the immediate geological past, should be deferred until it is too late to complete the record, by giving precedence altogether to the ancient formations. The rocks and fossils of the older formations will remain undisturbed for ages, as in the past, awaiting the leisure of the student of geology; while the delicate tracings of the latest pre-modern epochs are liable to fade away rapidly before the advancing settlement of the country. Nor can it be maintained that the processes that gave them birth, and which are still active in the formation of soils, are not scientifically, as well as practically, at least of equal interest with those that formed the older rocks. It is true that their study does not offer the easy rewards of the naming of new fossils, minerals, and rocks, which in times not yet belonging to the far past seemed to be the chief aim of students of geology; but they are none the less worthy of the highest scientific effort, and their practical results bear on products of an importance at least as great as those of the richest mines.

E. W. HILGARD.

Berkeley, Cal., May 1.

Queries.

32. HUMAN BEINGS AS PACK-ANIMALS.—In studying the history of transportation, I have ascertained that the first pack-animals were human beings, — men and women. Long before any of the animals were domesticated as beasts of burden, there were common carriers moving vast quantities of merchandise about the world. They toted (carried on the head); they hung great loads to their foreheads by means of a strap connected with a pack, wallet, basket, or frame on the back; they 'shouldered' their burdens, with or without yokes, front and rear, on one shoulder (like the Chinese) or on both shoulders (like the Dutch); they strapped their primitive knapsacks to their shoulders; they harnessed themselves to a load, as they did afterwards dogs, reindeer, horses, etc. Now, I should also like to know how much a man can tote, how much a woman can tote, and how long a time, without resting, the toting may go on. I should also like to know how much a man or a woman can carry in any particular manner, and how long a time the operation can be kept up without resting. The weight multiplied by the time will give a rough unit of human endurance. I shall be extremely obliged to any one who will give me valuable information on this subject.

O. T. MASON.

Washington, D.C., May 10.

Answers.

22. WASP-STINGS.—It is a fact not generally known, that, if one holds his breath, wasps, bees, and hornets can be handled with impunity. The skin becomes sting-proof, and holding the insect by the feet, and giving her full liberty of action, you can see her drive her weapon against the impenetrable surface with a force that lifts her body with every stroke; but, let the smallest quantity of air escape from the lungs, and the sting will penetrate at once. I have never seen an exception to this in twenty-five years' observation. I have taught young ladies with very delicate hands to astonish their friends by the performance of this feat; and I saw one so severely stung as to require the services of a physician, through laughing at a witty remark of her sister, forgetting that laughing required breath. For a theory in explanation, I am led to believe that holding the breath partially closes the pores of the skin. My experiments in that direction have not been exact enough to be of any scientific value, but I am satisfied that it very sensibly affects the amount of insensible perspiration.

W. L. WILDER.

Somerville, Mass., May 7.